

Scaling in Modulated Systems and Re-entrance of Order

O. Portmann, A. Vindigni, and D. Pescia

Laboratorium für Festkörperphysik, ETH Zürich, 8093 Zürich, Switzerland

(Dated: July 22, 2009)

We propose scaling invariance of d -dimensional modulated systems, in which modulation of order is produced by a frustrating long-ranged interaction decaying spatially with some power α of the inverse distance. Over the entire temperature range in which a modulation length is defined, measurable physical quantities acquire scaling exponents. One outcome of the scaling analysis is the existence, for $\alpha > d$, of an anomalous state in which the tendency of matter to order *increases* with temperature. We suggest that this anomaly is responsible for a re-entrance phenomenon in two-dimensional dipolar frustrated magnetic systems.

Modulation of order is a general motive in chemistry, biology, and physics [1, 2]. The modulated order parameter may represent quantities as diverse as the spin density [3, 4, 5], the charge density in any type of strongly correlated classical or quantum system [2, 6, 7, 8], the volume fraction of diblock copolymers, the concentration of amphiphilic molecules and other chemical species [2, 9], or dipolar bosons in an optical lattice [10]. However, modulated systems tend to show common characteristics such as the morphology of the various patterns and the occurrence of transitions among them [1], as in liquid crystals [11] or two-dimensional melting phenomena [12]. This tendency to common behavior indicates universal underlying principles [7, 8, 13].

In this Letter, we consider d -dimensional systems where the spatial modulation of the order parameter is produced by the competition of a short-ranged interaction favoring local order and a weak but long-ranged frustrating interaction decaying with some power α of the inverse distance. We propose that these systems are scaling invariant along the entire temperature range where some characteristic modulation length L exists. Accordingly, measurable physical quantities are characterized by scaling exponents which only depend on fundamental symmetries of the system, like d and α . One outcome of the scaling analysis is that modulation of order is only stable if α is less than an upper limit dependent on d . A further outcome is that, for $\alpha > d$, the tendency of the modulated system to order – measured by the compression modulus B – can *increase* with temperature. Among the systems that are predicted to show this anomaly are a model of two-dimensional Coulomb frustrated phase separation [8] and two-dimensional ferromagnets frustrated by the dipolar interaction [3, 4, 5]. We report experimental data on perpendicularly magnetized ultrathin Fe films on Cu(100) ($\alpha = 3 > d = 2$) [14, 15] that confirm the anomalous growth of B with temperature. This growth of B goes along with a re-entrance [16] of the more ordered stripe pattern [14].

Following Muratov [2], a universal effective Landau-Ginzburg-Wilson (LGW) Hamiltonian (per volume) $\mathcal{H}$ describing systems frustrated by a long-ranged interaction

reads [17]

$$\mathcal{H}[\phi(x), T] = \frac{1}{V_D} \int \left\{ \frac{J}{2} \phi_0^2 (\nabla \phi(x))^2 - T \phi_0^4 \left[\frac{\phi^2(x)}{6} - \frac{\phi^4(x)}{12} \right] \right\} d^D x + \frac{1}{V_D} \lambda \phi_0^2 \iint \phi(x) \phi(x') G_\lambda(|x-x'|) d^D x d^D x', \quad (1)$$

x being a vector in a D -dimensional space and V_D the volume of the system. Without the long-ranged interaction ($\lambda = 0$), this functional has macroscopic phase separation, with an order parameter ϕ_0 appearing below a critical temperature T_C . The field $\phi(x)$ is a scalar order parameter (rescaled with ϕ_0). The gradient term in the first line mimics the short-ranged exchange contribution. It is inaccurate for sharp walls at low temperatures, where a more suitable expression exists [4]. For large separations ($|x-x'|$), the kernel of the long-ranged interaction $G_\lambda(|x-x'|)$ is assumed to behave asymptotically like $|x-x'|^{-\alpha}$. $\alpha = 1$ corresponds to the Coulomb interaction and gives rise to the Coulomb Frustrated Ising Ferromagnet (CFIF) [6]. The kernel of the Dipolar Frustrated Ising Ferromagnet (DFIF) [20] has a more complicated structure but can also be treated with Eq. (1). We assume $\frac{J}{\lambda} \gg 1$ and hence a mono-dimensional modulation of length $L \gg 1$ along a coordinate x_1 in a system with a large extent in d dimensions. In the remaining $D-d$ dimensions, the system is assumed to have a finite thickness $\mathcal{O}(\delta) \ll L$.

At a given temperature T , we expect that a minor change of the modulation length L will have a limited effect on the order-parameter profile $\phi(x)$. We therefore postulate the invariance of $\phi(x)$ with respect to a scaling of all lengths by a constant u :

$$\phi(u \cdot x_1, u \cdot L) = \phi(x_1, L) \quad , \quad (2)$$

i.e. ϕ to be a function of one single variable $\bar{x}_1 \doteq \frac{x_1}{L}$; we thus introduce $\bar{\phi}(\frac{x_1}{L}) \doteq \phi(x_1, L)$. Considering only the asymptotic behavior of G_λ , we can now transfer the L -dependence of $\mathcal{H}$ to prefactors L^Δ with suitable scaling exponents Δ . Per unit volume, the part due to the exchange interaction scales like $\mathcal{O}(J \cdot \phi_0^2 \cdot L^{\Delta_J})$ and the part due to the long-ranged interaction like $\mathcal{O}(\lambda \cdot \phi_0^2 \cdot L^{\Delta_\lambda})$, with $\Delta_J = -1$ (sharp walls between adjacent domains) or -2

(extended walls) and $\Delta_\lambda = d - \alpha$. Some results of this scaling hypothesis are summarized in Table I and will be discussed in the following.

– The generalized Hellmann-Feynman theorem [18] equates the derivative of the Helmholtz free energy $\mathcal{F}$ per volume with respect to the parameter L with the canonical average over all configurations $\bar{\phi}(\bar{x}_1)$ of the derivative of $\mathcal{H}$ with respect to L :

$$\frac{\partial \mathcal{F}(L, T)}{\partial L} \doteq \left\langle \frac{\partial}{\partial L} \mathcal{H}[\bar{\phi}(\bar{x}), L, T] \right\rangle_{\bar{\phi}(\bar{x})}. \quad (3)$$

The characteristic modulation length $L_\lambda(T)$ is defined as the solution of the equation $\frac{\partial \mathcal{F}(L, T)}{\partial L} = 0$ and, when finite, scales like $(J/\lambda)^{\Delta_{L_\lambda}}$ with $\Delta_{L_\lambda} = 1/(\Delta_\lambda - \Delta_J)$ (third and fourth columns). Note that, in general, walls are very thin at low temperatures ($\Delta_J = -1$, third column) and broaden when the temperature is increased ($\Delta_J = -2$, fourth column) [20]. This produces the cross-over of the exponent Δ_{L_λ} observed in the third and fourth columns. Accordingly, scaling predicts a decrease of the equilibrium modulation length with temperature. Particularly striking is the cross-over (and the corresponding decrease of $L_\lambda(T)$) for the $(D = 3, d = 2)$ -DFIF. For $\Delta_J = -1$ ($T \approx 0$), we have $\Delta_{L_\lambda} = \infty$, which suggests an exponential dependence on J/λ (confirmed by detailed computations [4, 8]; see Ref. [19] for $d = 1$ and $\alpha = 2$, which also give $\Delta_{L_\lambda} = \infty$). At higher temperatures, L_λ only depends linearly on J/λ . For $J/\lambda \gg 1$, this means a decrease of the modulation length by several orders of magnitude [20].

– The compression modulus $B_\lambda(T)$ [4, 21] (also called Young modulus [22]) measures the energy cost associated with deviations from the equilibrium modulation length $L_\lambda(T)$ and is given by

$$B_\lambda(T) \doteq L_\lambda^2 \cdot \frac{\partial^2 F(L, T)}{\partial L^2} \bigg|_{L_\lambda(T)}, \quad (4)$$

which, in virtue of the Hellmann-Feynman theorem, is equivalent to

$$L_\lambda^2 \left\langle \frac{\partial^2}{\partial L^2} \mathcal{H}[\bar{\phi}(\bar{x}), L, T] - \frac{1}{T} \left(\frac{\partial}{\partial L} \mathcal{H}[\bar{\phi}(\bar{x}), L, T] \right)^2 \right\rangle_{\bar{\phi}(\bar{x})} \bigg|_{L_\lambda}. \quad (5)$$

Positivity of B_λ is a necessary condition for the stability of the modulated state. The scaling analysis, applied to Eq. (5), shows that $\langle \frac{\partial^2}{\partial L^2} \mathcal{H} \rangle|_{L_\lambda} \sim \lambda(\lambda/J)^{\frac{2-\Delta_\lambda}{\Delta_\lambda-\Delta_J}}$ and $\frac{1}{T} \langle (\frac{\partial}{\partial L} \mathcal{H})^2 \rangle|_{L_\lambda} \sim (\lambda^2/T)(\lambda/J)^{\frac{2-2\Delta_\lambda}{\Delta_\lambda-\Delta_J}}$, so that positivity of B_λ in the limit $J/\lambda \gg 1$ requires

$$\alpha - d < -\Delta_J. \quad (6)$$

This result generalizes the one obtained in Ref. [13] for $T = 0$ and $d = 1$. When the modulated state is stable, $B_\lambda \approx \lambda \cdot \phi_0^2 \cdot L_\lambda^{\Delta_{B_\lambda}}$ (for Δ_{B_λ} , see fifth column). Note that

		$\Delta_{L_\lambda}^{\Delta_J=-1}$	$\Delta_{L_\lambda}^{\Delta_J=-2}$	$\Delta_{B_\lambda}^{\Delta_J=-1,-2}$
(D, d)	α	$\frac{1}{d-\alpha+1}$	$\frac{1}{d-\alpha+2}$	$d - \alpha$
DFIF(3,3)		$\frac{1}{2}$	$\frac{1}{3}$	1
DFIF(3,2)	3	∞	1	-1
CFIF(3,3)	1	$\frac{1}{3}$	$\frac{1}{4}$	2
CFIF(3,2)	1	$\frac{1}{2}$	$\frac{1}{3}$	1

TABLE I: Scaling exponents Δ_{L_λ} and Δ_{B_λ} for general d and α and for some special cases. The asymptotic long-range interaction of the $(D = 3, d = 3)$ -DFIF does not follow a simple $|x - x'|^{-\alpha}$ -law. It therefore cannot be assigned a value of α and the exponents have to be determined separately.

an anomalous L -dependence appears for $d - \alpha < 0$ (cf. the $(D = 3, d = 2)$ -DFIF). B_λ *increases* (by several orders of magnitude in particular cases) with decreasing L_λ . This class of matter exhibits a state of potentially anomalous compression modulus, in which it becomes more difficult for the system to accommodate disorder involving deviations from the equilibrium modulation length when the temperature is increased; the system becomes *less* responsive to fluctuations at higher temperatures.

The validity of Eq. (2) is corroborated by a number of findings.

– First, we have computed equilibrium profiles $\phi_{\text{eq}}(x, L)$ for the $(d = 2)$ -DFIF, within the Mean-Field Approximation (MFA) of the functional Eq. (1) [20, 23]. At a given temperature, the MFA finds profiles that minimize the functional at a given L (black and light curves in Fig. 1(a)). $L_\lambda(T)$ corresponds to the profile (black curve) that minimizes $\mathcal{H}[\phi_{\text{eq}}(x, L), T]$ with respect to L and is found to decrease with temperature (not shown [20]). All profiles virtually collapse onto one single “scaling profile” in Fig. 1(b) when plotted as a function of $\frac{x}{2L}$. Figure 1(c) shows that the numerical $B_\Omega(T)$ (black curve) for the $(d = 2)$ -DFIF virtually coincides with the scaling prediction (light curve). In addition, it shows the anomalous growth with T predicted by scaling. Figure 1(d) shows the almost perfect coincidence between the numerical and the scaling result for $B_Q(T)$ in the $(d = 3)$ -DFIF and, in the inset, the $(d = 2, 3)$ -CFIF ($\alpha = 1$). $B_Q(T)$ decreases smoothly with temperature and does not have any anomaly.

– Second, we point out that Eq. (2) is trivially fulfilled at $T = 0$, as the order parameter field consists of segments with length L and alternating values ± 1 . We have verified the stability of the scaling hypothesis and of the scaling dimensions at finite temperatures by introducing small deviations from the strict mono-dimensional order, modifying $\phi(x_1)$ to $\phi(x_1 - \epsilon_{hk} \cos(kx_2) \cos(hx_1))$. This replacement establishes an elementary excitation displacing domain walls along the x_1 -coordinate while bending them along the x_2 -direction and accounts for the experimental fact that stripes are never perfectly straight in real systems [24] at finite temperatures. Such




FIG. 1: (a): (color online). Profile of the order parameter in a $(d=2)$ -DFIF ($\delta=1$, $\Omega/J=0.08$) for different values of L ($0.7L_\Omega$, L_Ω and $1.3L_\Omega$). (b): Collapse of profiles onto one single curve in plot versus $\frac{x}{L}$. (c): Numerical $B_\Omega(T)$ for the $(d=2)$ -DFIF (black curve, $\delta=1$, $\Omega/J=0.08$) together with the scaling result (light line). (d): Numerical and scaling curves $B_\Omega(T)$ for the $(d=3)$ -DFIF ($\delta=200$, $\Omega/J=0.08$) and the $(d=2,3)$ -CFIF (inset, $\delta=1, 200$, $Q/J=0.0001$). Note that at low temperatures the wall is very thin and the numerical results are unstable. Light circles on the vertical axis indicate the values of B_λ at $T=0$ obtained analytically.

excitations are responsible e.g. for the Landau-Peierls instability that prevents order in $D=3$ for $d=1, 2$ [2, 3, 4]. Averaging the perturbed functional $\mathcal{H}[\phi(x_1 - \epsilon_{hk} \cos(kx_2) \cdot \cos(hx_1)), L, T]$ expanded up to order $(\epsilon_{hk})^2$ and powers $(h \cdot k)^n$ with $n \leq 4$ using the equipartition theorem $\langle \epsilon_{hk}^2 \rangle \approx \frac{T}{Jq^2 h^2 + \lambda q^{\alpha-3} k^4}$ [4] and summing over $(h, k) \leq 1/L$, we recover an effective mono-dimensional functional in which the exchange energy and the energy of the long-ranged interaction have, to leading order, the same scaling behavior, albeit with modified coupling constants $J \mapsto J + \mathcal{O}(T)$ and $\lambda \mapsto \lambda + \mathcal{O}(\lambda \frac{T}{J})$. This shows that scaling is conserved at finite temperatures and that the




FIG. 2: Experimental $B_\Omega(T)$ for an ultrathin Fe film on Cu(001) obtained by dividing $\phi_0(T)$ by $L_\Omega(T)$. The solid line results from fits to $\phi_0(T)$ and $L_\Omega(T)$ respectively and is a guide to the eye. The temperature range with stripe domains is shaded, the dark shade indicating mobile stripes. The two insets show SEMPA (Scanning Electron Microscopy with Polarization Analysis) images of the labyrinthine and the stripe pattern. The length of the white bar is $10 \mu\text{m}$, the thickness of the film 1.95 ML.

scaling dimensions are not changed by small perturbations. In particular, the Landau-Peierls instability destroys long-range positional stripe order but the loss of order proceeds in such a way that the modulation length and the compression modulus B_λ remain well-defined quantities also at finite temperatures [25].

– Third, the compression modulus B_Ω of perpendicularly magnetized ultrathin Fe films on Cu(100) [14] shows the anomalous enhancement with temperature predicted by scaling. In these systems, a strong magnetic anisotropy aligns the spins perpendicular to the film plane. Accordingly, the exchange interaction is frustrated by the (isotropic part) of the long-range dipolar interaction, i.e. it is a $(d=2)$ -DFIF and has a characteristic modulation length [1, 14], up to a temperature T_C at which ϕ_0 vanishes. The data points in Fig. 2 display $B_\Omega(T)$ as determined experimentally using the scaling result. The solid curve is obtained from fits to $\phi_0(T)$ and $L_\Omega(T)$. The decrease of $\phi_0(T)$ and the appearance of mobile stripes just below T_C makes the experimental determination of $B_\Omega(T)$ difficult. Note that in zero applied magnetic field, the low-temperature state is striped, but transforms upon temperature increase into a more disordered labyrinthine pattern, displayed left in Fig. 2 [1, 14]. This transformation has also been reported in a micrometer-thick film, a $(d=3)$ -DFIF [1]. What is new in the ultrathin limit is that the labyrinthine pattern transforms back into the more ordered stripe pattern (pictured right in Fig. 2) when the temperature is increased. This re-entrance of order proceeds in the temperature region (light shaded area) where the widespread local configurations in the labyrinthine pattern with deviations from the equilibrium stripe width are increasingly

penalized by the increasing $B_\Omega(T)$. This suggests that the unusual re-entrance of stripe order is a response of the system to the extra elastic energy introduced by the increase of $B_\Omega(T)$. Note that the experimental temperature dependence of the stripe width [15] is smooth in the temperature range where the labyrinth-to-stripe transformation proceeds, which confirms the predicted insensitivity of L_Ω to deviations from mono-dimensional order. In the vicinity of T_C (dark shaded area), the stripes become mobile [15]. This mobility coincides with the range of temperatures where $B_\Omega(T)$ is expected to decrease and, accordingly, can no longer counteract thermal excitations which destroy stripe order.

In conclusion, we have presented a scaling procedure that is generally applicable to any pattern-forming system and allows re-writing the Hamiltonian (1) as a functional of a scaling-invariant scalar field and rescaled coupling constants $J \mapsto JL^{\Delta_J}$ and $\lambda \mapsto \lambda \cdot L^{\Delta_\lambda}$. The temperature dependence of L_λ , which may vary by several orders of magnitude, is thus expected to propagate in a non-trivial way to any thermodynamic observable. Among other possible implications, scaling predicts a yet undetected state of matter with a compression modulus increasing with temperature. This anomaly provides a simple explanation for an anomalous sequence of patterns observed in ultrathin ferromagnetic films. We show that a necessary condition for this anomaly is $\alpha > d$, which is fulfilled by the ultrathin limit of dipolar frustrated Ising ferromagnets (or equivalent models, see Ref. [8]), but not by dipolar frustrated films that are not strictly two-dimensional or in Coulomb frustrated systems. This explains why, despite the huge amount of experimental and theoretical work, no other instances of re-entrance have ever been observed in the many other pattern-forming systems [1]. So far, predictions and studies of topological transitions [12], although extensive, have neglected possible anomalous behavior of the compression modulus. Our analysis indicates the need to refine the various scenarios for topological phase transitions. In addition, we anticipate that tuning B_λ with an appropriate choice of temperature, thickness – and probably applied magnetic field – might provide new ways of producing two-dimensional melting transitions and allow investigating their elusive topological character as well as the controlled cross-over to more conventional three-dimensional melting. We also anticipate that other re-entrance phenomena – e.g. in an applied magnetic field – might arise because of the anomalous compression modulus.

We acknowledge the financial support of ETH Zurich and the Swiss National Science Foundation as well as fruitful discussions with P. Politi.

[1] M. Seul and D. Andelman, *Science* **267**, 476 (1995) and references therein.

- [2] C. B. Muratov, *Phys. Rev. E* **66**, 066108 (2002) and references therein.
- [3] T. Garel and S. Doniach, *Phys. Rev. B* **26**, 325 (1982).
- [4] A. B. Kashuba and V. L. Pokrovsky, *Phys. Rev. B* **48**, 10335 (1993); Ar. Abanov, V. Kalatsky, V. L. Pokrovsky, and W. M. Saslow, *Phys. Rev. B* **51**, 1023 (1995).
- [5] Kwok-On Ng and D. Vanderbilt, *Phys. Rev. B* **52**, 2177 (1995); A. D. Stoycheva and S. J. Singer, *Phys. Rev. Lett.* **84**, 4657 (2000); K. De'Bell, A. B. MacIsaac, and J. P. Whitehead, *Rev. Mod. Phys.* **72**, 225 (2000); D. G. Barci and D. A. Stariolo, *Phys. Rev. Lett.* **98**, 200604 (2007); S. A. Pighin and S. A. Cannas, *Phys. Rev. B* **75**, 224433 (2007) and references therein.
- [6] M. Grousson, G. Tarjus, and P. Viot, *Phys. Rev. E* **62**, 7781 (2000); **64**, 036109 (2001); M. Grousson, V. Krakoviack, G. Tarjus, and P. Viot, *Phys. Rev. E* **66**, 026126 (2002).
- [7] Z. Nussinov, J. Rudnick, S. A. Kivelson, and L. N. Chayes, *Phys. Rev. Lett.* **83**, 472 (1999) and Z. Nussinov, *ArXiv:cond-mat/0506554*.
- [8] R. Jamei, S. Kivelson, and B. Spivak, *Phys. Rev. Lett.* **94**, 056805 (2005); C. Ortix, J. Lorenzana, and C. Di Castro, *Phys. Rev. Lett.* **100**, 246402 (2008).
- [9] H. M. McConnell, *Annu. Rev. Phys. Chem.* **42**, 171 (1991).
- [10] C. Menotti, C. Trefzger, and M. Lewenstein, *Phys. Rev. Lett.* **98**, 235301 (2007).
- [11] C. Harrison et al., *Phys. Rev. E* **66**, 011706 (2002) and references therein.
- [12] K. J. Strandburg, *Rev. Mod. Phys.* **60**, 161 (1988) and references therein.
- [13] A. Giuliani, J. L. Lebowitz, and E. H. Lieb, *Phys. Rev. B* **74**, 064420 (2006); M. Biskup, L. Chayes, and S. A. Kivelson, *Comm. Math. Phys.* **274**, 217 (2007).
- [14] O. Portmann, A. Vaterlaus, and D. Pescia, *Nature* **422**, 701 (2003).
- [15] O. Portmann, A. Vaterlaus, and D. Pescia, *Phys. Rev. Lett.* **96**, 047212 (2006).
- [16] T. Narayanan and A. Kumar, *Physics Reports* **249**, 135 (1994); L. Greer, *Nature* **404**, 134 (2000); N. Schupper and N. M. Shnerb, *Phys. Rev. E* **72**, 046107 (2005).
- [17] The coupling constant λ amounts to $\Omega \doteq \mu_0(g\mu_B S)^2 \cdot (4\pi a^3)^{-1}$ in the dipolar case and to $Q \doteq q^2 \cdot (4\pi\epsilon_0 a)^{-1}$ for the Coulomb interaction respectively. All lengths in Eq. (1) are in units of the lattice constant a . For convenience, we set $k_B = 1$.
- [18] L. D. Landau and E. M. Lifshitz, *Statistical Physics*, Pergamon Press, 1980, p. 49.
- [19] E. Nielsen, R. N. Bhatt, and D. A. Huse, *Phys. Rev. B* **77**, 054432 (2008).
- [20] A. Vindigni, N. Saratz, O. Portmann, D. Pescia, and P. Politi, *Phys. Rev. B* **77**, 092414 (2008).
- [21] D. Sornette, *J. Physique* **48**, 151 (1987).
- [22] M. Kleman and O. D. Lavrentovich, *Soft Matter Physics*, Springer-Verlag, 2003, p. 144.
- [23] In order to obtain more precise MFA results at lower temperatures, we use the Bragg-Williams approximation for the entropy (cf. Ref. [20]) in Eq. (1).
- [24] R. de Koker, W. Jiang, and H. M. McConnell, *J. Phys. Chem.* **99**, 6251 (1995).
- [25] F. Cinti, O. Portmann, D. Pescia, and A. Vindigni, *Phys. Rev. B* **79**, 214434 (2009).